

Diagnostic Submission Guidelines: Respiratory Disease

The ideal samples for investigation of respiratory diseases are those that are collected within 4-8 hours of death, however, any sample is better than none, and the lab does not charge based on the number of samples submitted. **Both** fresh and formalin fixed tissues from **2-4** different sections of lung including the junction between normal and diseased tissue as well as the following samples should be included:

Organ	Formalin*	Fresh	Size Fresh Samples
Heart	Yes	Yes	4 x 4 x 2 cm
Lung (2-4 sections)	Yes	Yes	4 x 4 x 2 cm
Trachea	Yes	Yes	4 x 4 x 2 cm
Bronchial Lymph Node	Yes	Yes	4 x 4 x 2 cm
Mesenteric Lymph Node	Yes	Yes	
lleum	Yes	Yes	10 - 20 cm length

* A 2 x 2 x 1 cm piece of tissue from the organs listed above should be placed in formalin. The ratio should be 9 parts formalin to 1 part tissue for proper fixation. The WVDL recommends using a 16 ounce wide mouthed jar that contains 8 ounces of 10% buffered formalin (Fischer catalog # 23-426-797). Since the formalin jars tend to leak; **please place the formalin jar inside a Ziploc**® **bag**.

Shipping Requirements

- Veterinarians should completely fill out the WVDL General Submission Form which includes the **age** of the animal. An electronic copy is available at <u>www.wvdl.wisc.edu</u>. Click on the forms link to download the submission form.
- The laboratory should receive the samples within 24-36 hours so samples collected on Friday should be refrigerated and shipped on Monday.
- Package the samples with a sufficient number of ice packs to ensure they remain **cold** during shipment.
- Check out our special shipping rates here.
- **WINTER SHIPMENTS:** Please add isopropyl alcohol (1:10 ratio) to the formalin for formalin fixed tissues to prevent freezing during shipment.